
A Summary of ACPI Climate Downscaling Studies for the Western United States

L. Ruby Leung, Yun Qian, and Xindi Bian
Pacific Northwest National Laboratory, Richland, WA
John Roads, Jongil Han, and Simon Mason
Scripps Institute of Oceanography, San Diego, CA

2001 AGU Fall Meeting December 10-14, 2001, San Francisco, CA

Objectives

- To evaluate downscaling methods for the western U.S.
- To intercompare dynamical and statistical downscaling methods
- To provide downscaled climate scenarios for climate impacts assessment





Climate of Western U.S. is Strongly Affected by Terrain:

Model capability to resolve terrain features is critical


Observed snow pack in March, 1998

Observed mean annual precipitation





ACPI Downscaling Results



- Two regional models (MM5, RSM)
 have been evaluated when driven
 by realistic large scale conditions
 from the NCEP/NCAR and
 ECMWF reanalyses.
- Intercomparison of regional simulations have been performed at the Columbia River and Sacramento-San Joaquin basins.
- Both MM5 and RSM are being used to downscale the PCM control and future climate conditions.
- Regional simulations have been provided to impact assessment team.

Data Archives

• PCM outputs:

6-Hourly (5.5 GB); Daily (0.4 GB); Monthly (0.3 GB) – total 2.2 TB for 350 years and selected hourly data


• MM5 outputs:

Daily (0.2 GB), 3-Hourly (0.8 GB), 6-Hourly (6.4 GB) – total 0.8 TB for 110 years


RSM outputs:

3-Hourly (1.0 GB) – total of 75 GB for 75 years


Seasonal Precipitation: Simulation Driven by ECMWF Reanalyses




Seasonal Precipitation: Simulation Driven by NCEP/NCAR Reanalyses




Comparison of Extreme Precipitation





Basin Mean Monthly Precipitation




Comparison of Simulations












Observed and Simulated El Nino Precipitation Anomaly



PCM Simulated Realistic Seasonal Cycle of Precipitation and Temperature




PCM Annual Temperature and Precipitation at the Columbia River Basin




ControlBAU 1BAU 2BAU 3


Seasonal Precipitation: PCM and RCM Control Simulations




Ensemble Mean PCM Temperature Signal




Ensemble Mean PCM Precipitation Signal




Seasonal Temperature: climate Signal




Seasonal Temperature: climate Signal




Seasonal Precipitation: Climate Signal




Seasonal Precipitation: Climate Signal





Monthly Temperature: Climate Signal Simulated by PCM and RCM





Seasonal Precipitation: Climate Signal Simulated by PCM and RCM





Conclusions

- Differences exist between MM5 and RSM simulations for both summer and winter seasons.
- Interannual variability is realistically simulated by both models at the river basins when driven by reanalyses.
- PCM control simulation reproduces seasonal cycle of temperature and precipitation at the river basins.
- Ensemble means over 20 years are more similar to each other than 10-year averages.
- Downscaling accentuates temperature and precipitation (especially extreme) changes over mountainous regions.